

GABA_A receptor plasticity in neuropathic pain: pain and memory effects in adult female rats

Azeez O. Ishola^{a,c}, Adesola T. Ademola^a, Rosemary K. Allen^a, Babafemi J. Laoye^b, Oluwamolakun T. Bankole^b, Moyosore S. Ajao^c, Philip A. Adeniyi^a

^aNeural Toxicity Unit, Anatomy Department, Afe Babalola University Ado-Ekiti, ^bBiological Science Department, Afe Babalola University Ado-Ekiti, ^cAnatomy Department, University of Ilorin, Ilorin, Nigeria

Correspondence to Azeez O. Ishola, BSc, MSc, Anatomy Department, Afe Babalola University Ado-Ekiti, Nigeria. PMB 5454. Zip Code: 360211. e-mail: ao.ishola@abuad.edu.ng

Received 9 May 2018

Accepted 10 February 2019

Egyptian Pharmaceutical Journal 2019, 18:8–15

Background

Neuropathic pain has been shown to increase excitability of neurons. This indicates altered inhibitory mechanism of the nervous system.

Objective

This work was aimed to assess GABA_A receptors plasticity in the brain and spinal cord.

Materials and methods

Fifteen adult female rats were used. Ten animals have their sciatic nerve ligated with no treatment (LIG), and with diazepam treatment for 14 days (LIG+GABA) and the other five were used as the sham group. Pain was assessed using a hot plate and formalin test, while the spatial memory was assessed using Y-maze. At the end of the treatment, the animals were euthanized and fixed using the transcardial perfusion fixation method. The spinal cord, cingulate cortex, and the hippocampus were serially sectioned and stained for GABA_A receptor immunohistochemically. Quantification was done using ImageJ software. Data were analyzed using one-way analysis of variance and Newman Tukey post-hoc test significant level was set at *P* less than 0.05.

Results

A low level of pain was observed in LIG and LIG+GABA animals on both formalin and hot plate test compared with the control. Memory impairment was found only in the LIG+GABA group. Stereology counting showed that GABA_A receptors reduced in the dentate gyrus of the hippocampus of LIG-treated animals which was reversed in LIG+GABA, but in the cingulate cortex, GABA_A receptors were increased in LIG animals and LIG+GABA more than the control while the spinal cord shows no significant difference.

Conclusion

GABA_A agonist treatment did not alleviate the symptoms of neuropathic pain due to GABA signaling changing to excitatory in nature.

Keywords:

cingulate cortex, GABA_A receptor, hippocampus, neuropathic pain, spinal cord

Egypt Pharmaceut J 18:8–15

© 2019 Egyptian Pharmaceutical Journal
1687-4315

Introduction

Neuropathic pain (NP) is known to arise due to injury or dysfunction in the peripheral and the central nervous system [1–3]. This dysfunction in the pain pathway leads to hypersensitivity (i.e. the organism feels pain with no stimuli), allodynia (i.e. painless tactile stimulus or warmth elicit pain sensation), and hyperalgesia (i.e. painful stimuli elicits greater intensity of pain sensation) [3]. Sometimes NP is chronic and can results from diabetes, chemotherapy, amputation, and others [4] and it can be induced by chronic constriction injury (CCI) in laboratory animals [5,6] by ligating the spinal cord or peripheral nerve fiber [7]. Using this method, it is discovered that injury to the nerve fibers causes hyper-depolarization in the surrounding spared fiber which are responsible for the chronic pain

experienced in NP [8–12]. Another mechanism that has been reported is the disinhibition of the pain pathway and loss of GABA neurons [13,14].

GABA/glycine inhibitory neurons in the dorsal horn are usually activated to reduce pain transmission [3]. With the loss of inhibition, excitotoxicity in NP may lead to the loss of these neurons leading to hyperalgesia [15]. Previous reports have indicated that there are plastic changes in the central nervous system due to CCI [8,16–18]. These changes are responsible for the brain

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

autoregulatory mechanism to adapt to the injury [19], leading to further propagate the pain experienced [20]. Most of these changes induce the loss of GABA neurons in the spinal cord (dorsal horn) and cingulate cortex (area responsible for sensory processing in rodents) [13,14].

Memory decline is another symptom associated with NP [15,21]. Although the mechanism underlying the decline in memory is not yet understood, some reports have shown that this may be due to excitotoxicity [15]. Mutso *et al.* [22] reported hippocampal-mediated behavioral changes in NP which they associate to ipsilateral upregulation of dpErk molecules.

This study was designed to study whether NP induces GABA_A receptor plasticity in the spinal cord, cingulate cortex, and the hippocampus and activating the receptor will ameliorate the pain and memory deficit in NP.

Materials and methods

Experimental animals

Fifteen adult female rats with an average weight of 150 g were procured from the animal holdings of the Department of Anatomy, Afe Babalola University Ado-Ekiti, Nigeria. The rats were housed in the standard plastic cages of five animals per cage. Food (ret pellets) and water were provided *ad libitum*.

Ethical statement

This experiment was carried out in accordance with the Nigerian National Ethical Code on animal research with formal approval from Afe Babalola University Ethics Committee.

Drug preparation

Commercially available GABA_A receptor agonist (diazepam) injection was purchased from RichyGod International Ltd (Lagos, Nigeria). The drug was prepared into solution by dissolving the ampules in normal saline before administering it to the animals.

Animal grouping

The animals were divided into three groups (sham, ligated, and ligated with treatment of five animals each). Before sciatic nerve ligation, all animals were anesthetized using ketamine Sham (SHAM): animals had their right sciatic nerve exposed and closed up back without ligation. The animals later received 2 ml/kg body weight (BW) normal saline.

Ligated (LIG) group: animals had their right sciatic nerve ligated and later received 2 ml/kg BW normal saline.

Ligated with treatment (LIG+GABA) group: animals had their right sciatic nerve ligated and later treated with 10 mg/kg BW of GABA_A receptor agonist (diazepam).

All administrations were done intraperitoneally and daily for 14 days.

Sciatic nerve ligation procedure

NP was induced by sciatic nerve ligation using the Bennett and Xie model [7]. The animals were sedated with ketamine (2 ml/kg BW intraperitoneal). The animals were placed in a pronated position when deeply anesthetized and were immobilized to the surgical table with a clip. The skin on the right thigh region was cleaned with a cotton wool soaked in ethanol on the dorsal part. The incision was made in this region to expose the sciatic nerve at the mid-thigh level; the nerve was separated from the surrounding tissues, it was raised up with forceps, and three ligatures were carefully tied around the sciatic nerve with 6-0 silk surgical sutures – VCP 496. P30 [23].

Animals in the sham control group had their thighs opened to expose the sciatic nerve and their thigh sutured back without ligating the sciatic nerve.

After the surgical procedure the incised region was treated with procaine penicillin (antibiotics) to avoid infection. Treatment was started 3 days after the surgical procedure.

Behavioral studies

The animals were exposed to a battery of tests to assess the level of pain and memory due to sciatic nerve ligation and or diazepam treatment.

Hot plate test

The aim of this test was to determine the level of pain sensation in the animals termed thermal hyperalgesia [24]. A transparent box made of Pyrex was placed on the hot plate to prevent the animals from roaming around on the hot plate and the animal was placed within the box on the regulated hot plate set at 55°C. Timing started when the animals were placed in the box. Once the animal starts flicking/licking its paws or tail, the timer was stopped and the time was recorded. After this, the animal is returned back to its home cage.

Formalin test

This involves injecting the paws of the animal with 2% formal saline [25]. The rat was then placed down and the number of times it beats (taps) its leg to the ground

in 1 min is taken and recorded as an acute stage. Twenty minutes later, the same recording was taken again from the animals termed the chronic phase which was centrally mediated.

Y-maze test

The Y-maze was used in assessing the spatial working memory of the rats [26]. The rat was placed facing the edge of the maze and monitored on a screen while being timed for 5 min. Visiting the three different arms consecutively was termed right decision (right) while visiting one arm more than once in three alternations was termed wrong decision (wrong). Memory index was calculated as the percentage of right decisions for each animal.

The behavioral studies were done in the order of Y-maze, hot plate, and formalin tests with a day interval each.

Animal sacrifice

After the last behavioral protocol, the animals were deeply anaesthetized with 10 ml/kg body weight of ketamine intraperitoneally, after which the animals were fixed transcardially by flushing the blood with 0.9% normal saline and later with 10% formal saline solution. The brain and the spinal cord were dissected out and postfixed in 10% formal saline solution.

Detection of GABA_A receptor in the spinal cord, cingulate cortex, and the hippocampus using immunohistochemical staining

Immunohistochemical staining was performed using the heat method of antigen retrieval. Brain and spinal cord slices already processed and paraffin embedded were sectioned serially, placing every 10 sections on the slides. Slides were baked in the oven at 50°C for 30 min. Then the slides were placed in xylene for 10 min, rehydrated in 100, 90, 70, and 50% of alcohol for 10 min each and then rinsed with distilled water. The slides were placed in hot citric acid solution (antigen retrieval solution) at a pH of 7.0 heated to 70°C for 40 min till it cools down to room temperature. Then the slides were rinsed with distilled water and 1xPBS thrice for 5 min each.

Tissue area was encircled using a PAP pen and then incubated in H₂O₂ for 30 min at room temperature to block endogenous peroxidase followed by protein block solution for 10 min at room temperature and the slides were rinsed with PBS in between the incubation.

The slides were incubated in rabbit anti-GABA_A receptor polyclonal antibody from Novus Biologicals

(1:100 NB100-61096, Novus Biologicals Centennial CO, USA) at room temperature for 180 min. Later they were incubated with goat polyvalent antimouse/rabbit secondary antibody from Abcam (1:100 ab64258, Abcam Cambridge, MA, USA) at room temperature for 60 min. Chromogen development was done in accordance with the manufacturer manual of DAB Substrate kit (ab64238; Abcam Cambridge, MA, USA). The slides were counterstained in 1% aqueous hematoxylin solution from Emsdiasum (26042-1; Emsdiasum Hatfield, PA, USA) for 20 min and dehydrated back in 50, 70, 90, and 100% of alcohol for 10 min each and later were cleared in xylene for 10 min and were covered slipped with DPX and then left to dry and viewed under a microscope (Olympus microscope attached with Winjoe 5MP cameroscope).

Stereology

The region of interest [dorsal horn of the spinal cord, anterior cingulate cortex, and dentate gyrus (DG) of the hippocampus] was viewed under a microscope at x40 objective. Positive cells were counted using ImgaeJ software (ImageJ Wisconsin, USA) and recorded.

Statistical analysis

Data were expressed using mean±SEM and were analyzed using one-way analysis of variance. Newman Tukey post-hoc test was done when the analysis of variance shows significance. The *P* value was set at 0.05. The analysis was done using GraphPad Prism software (San Diego, CA, USA).


Results

Increased hyperalgesia seen in LIG and LIG+GABA

Diazepam treatment fails to ameliorate pain perception in ligated animals as seen from the hot plate test. Ligated animals (LIG) and animals treated with diazepam (LIG+GABA) showed a significant reduction in the time spent before the experience of pain compared with SHAM animals (Fig. 1a). No significant difference was seen between the GABA-treated and ligated animals.


Paw flinching during the formalin test showed that the LIG+GABA-treated group exhibited the highest number of flinches both at an acute and chronic phase which is statistically significant compared with the SHAM group. During the acute phase, no significant difference was observed between the LIG group and SHAM group as this pain is mediated peripherally contrary to the chronic phase which was

Figure 1



The graph showed the results of the pain test. (a) The latency of pain in animals during hot plate test set at 55°C. The ligated animals (LIG, LIG+GABA) had a low threshold of pain as seen from the graph with a significant decrease ($P < 0.05$) in the latency of time. Animals treated with GABA agonist after ligation (LIG+GABA) did not show improvement in their threshold of pains compared with the SHAM and no significant difference was observed in the pain threshold of ligated and GABA-treated ligated animals. (b) The pain perception results from formalin test (acute and chronic phase). At acute phase animals treated with GABA agonist (LIG+GABA) showed the highest number of paw flinches which was significantly higher ($P < 0.05$) compared with the sham and ligated animals only (SHAM, LIG). No significant difference was observed when SHAM and LIG were compared. Ligated animals and those treated with GABA agonist (LIG, LIG+GABA) have increased the number of flinches, which are significantly higher ($P < 0.01$) compared with sham animals. A significant difference was not observed when LIG and LIG+GABA when compared in the chronic phase.

Figure 2



Graph showing the spatial memory index of animals in Y-maze test. No significant difference was observed between the sham animals and the ligated animals (SHAM vs. LIG). Animals treated with GABA agonist after ligation of the sciatic nerve (LIG+GABA) showed a significant reduction in their spatial memory index compared with the sham animals and ligated animals.

mediated centrally where LIG is significantly differenced from the SHAM group but not to the LIG+GABA-treated group (Fig. 1b).

Neuropathic pain and spatial memory

Spatial memory impairment was observed only in the LIG+GABA group which was significantly lower compared with SHAM and LIG groups. Moreover, no significant difference was observed between SHAM and LIG (Fig. 2).

NP effect on hippocampal, cingulate cortex, and spinal cord GABA_A receptor plasticity

GABA_A receptors were expressed at all levels of the spinal cord of experimental animals (Fig. 3a). Stereological counts of positive cells in the spinal cord showed that the sciatic nerve ligation slightly reduced the expression of GABA receptors with further reduction in GABA receptor agonist treatment (Fig. 3b).


Hippocampal and cingulate serial sections expressed GABA_A receptors (Fig. 4a). Stereology counting showed that the receptors were reduced in the DG of the hippocampus of ligated animals which was reversed by GABA_A receptor agonist treatment (LIG+GABA) more than the control (Fig. 4b). Cingulate cortex (Cg) which processes sensory inputs in rodents expressed more GABA receptors in ligated animals and further increased by GABA_A receptor agonist treatment (LIG, LIG+GABA) more than the control (Fig. 4b).

Discussion

Pain perception

In this study, formalin and hot plate test were used to monitor the pain response in the animals. Hot plate and formalin test showed that the ligated animals (animals induced with NP) showed hyperalgesia because they responded to the stimulus faster than control animals (sham), and the treatment with GABA_A receptor agonist (diazepam) did not

Figure 3




GABA receptor plasticity in the spinal cord at all levels. (a) Immunohistochemical slides of the spinal cord of experimental animals stained for GABA receptors at $\times 40$. GABA receptors were expressed in both anterior and posterior horns of the spinal cord at all levels. (b) GABA receptor immunopositive cell counts in the posterior horn of the spinal cord of experimental animals. There was a steady decline in the number of expression in the ligated animals and those treated with GABA receptor agonist (LIG, LIG+GABA) compared with the control.

alleviate hyperalgesia in the animals. Previous studies have shown that the sciatic nerve ligation model of NP induced loss of activity on the injured nerve and hyperexcitation of the surrounding nerve fibers [15,27]. Hyperexcitation of the nerve fibers has been involved in the pathophysiology of NP [27–29]. It has been suggested that this is due to the system plasticity to correct for the activity of the injured nerve [19]. In a report by Abdulmajeed *et al.* [15], peripheral and center nerve damage was reported, which was associated with excitotoxicity due to sciatic nerve ligation of NP.

GABA neurons in the spinal cord are inhibitory in nature; they regulated the level of pain transmission through the spinal cord. Activating GABA_A receptors (using diazepam) failed to improve pain perception as it further increased pain perception in this model of NP. This was similar to what was reported by Ran *et al.* [28], who showed that muscimol topically injected at the site of nerve damage did not alleviate thermal hyperalgesia in NP rats. Although, in a report by Chen *et al.* [30], they showed that single intraperitoneal injection of diazepam at 9 days after CCI alleviates mechanical allodynia and thermal

Figure 4



GABA receptor plasticity in the hippocampus and cingulate cortex. (a) Slides show the dentate gyrus of the hippocampus and cingulate cortex of experimental animals stained for GABA receptors at $\times 100$. GABA receptors were expressed in all regions under view. (b) GABA receptor immunopositive cell counts in the dentate gyrus of the hippocampus (DG) and cingulate cortex (Cg) expressed as a percentage of control. GABA receptor expressing cells were reduced in the dentate gyrus of the hippocampus of ligated animals, whereas it is more in the GABA-treated ligated animals. Cingulate cortex where sensory processing takes place had more GABA receptors expressing cells in ligated and GABA-treated ligated animals compared with control animals.

hyperalgesia in NP rats. This result was not consistent with other reports that showed that glycine which was a major inhibitory neurotransmitter in the spinal cord showed abnormal signaling and reduced expression in the chronic constriction model NP [29].

Another mechanism that explained the activation of GABA_A receptors not alleviating mechanical pain is that they may contribute to enhancing excitation and not inhibition. Ran *et al.* [28] reported that GABA_A

receptors of the adjacent intact dorsal root ganglion may be crucial in the development of hyperalgesia in NP. Ford *et al.* [31] reported that KCC2 signaling responsible for polarization for the inhibitory mechanism of GABA was impaired in NP. The authors also showed that administering adenosine receptor agonist did not have any analgesic effect until when KCC2 signaling is restored. This showed that the expression of KCC2 is altered in NP, which may mimic what happens during neurodevelopment

where NKCC1 was expressed more making GABA transmission to be excitatory [32].

Memory

Memory impairment has been associated with the development of NP [21]. The underlying mechanism associated with these has been proposed to be abnormal signal transmission and plastic changes due to the nerve injury [19]. Hippocampus the main structure responsible for memory formation is also reported to be involved in the transition from acute to chronic pain [33]. NP is reported to cause plastic changes in different regions of the brain with hippocampus inclusive [22,33,34]. The present study showed that ligated animals have no memory impairment on Y-maze compared with the control, but diazepam intervention leads to memory decline. Although memory deficit was reported in neuropathic animals no memory deficit was seen on Y-maze test [15].

Activation of GABA_A receptors leads to memory decline in NP in this study. NP has been shown to increase the level of GABA in the hippocampus [33]. Increasing GABA will stimulate the receptors in the brain, additional stimulation of GABA receptors led to hyperstimulation of this receptor which can lead to long-term depression leading to memory loss [35,36].

GABA_A receptor plasticity

NP has been reported to induce plastic changes in the spinal cord [8]. The major prone areas of these changes have been the dorsal horn since it carries the sensory signals to the brain [37]. Spinal cord inhibitory neurons played a regulatory role in dampening the hyperexcitation seen in NP has been shown to be affected by the plastic changes in NP [8,38]. The present study showed that sciatic nerve ligation induced a slight decrease in the number of GABA_A receptor expressing cells, which is further reduced with diazepam treatment. Although GABAergic neurons have been reported to reduce in the dorsal horn of the spinal cord [8,16–18], the receptors for the neurotransmitter released by these neurons also showed signs of reduction in correlation with production reduction. Although there was no significant difference in the loss of GABA_A receptors, this was similar to what was reported by Polgár and Todd [38].

Cingulate cortex which is responsible for sensory processing in rodents [2,39] is also affected by NP [14]. This region is a key component of the pain pathway. Plastic changes in this region have been hypothesized to be a key in the persistent pain experienced in NP [40,41]. The cortical interneuron reorganization has been identified to occur in the

cingulate cortex due to NP [14]. Loss of GABA-inhibitory transmission is known to be a key in the pathophysiology of NP [42]. There was an increase in GABA_A receptor expression in NP animals, activation of GABA_A receptor in the NP animals led to a further increased expression of GABA_A receptors. This may be due to the receptor dynamics to mop up the excess agonist from the system. The animals treated with GABA_A receptor agonist still have hyperalgesia. This indicated that increased GABA_A receptor activation was not leading to inhibition but more excitation of pain pathway as it was reported by Blom *et al.* [14], who showed that cingulate cortex disinhibition led to increased symptoms of NP.

GABA_A receptor was reduced in the DG of the hippocampus in neuropathic animals with increased expression in GABA_A receptor agonist-treated animals. The reduced expression of GABA_A receptor in neuropathic animals explained why there was no memory impairment at this stage on Y-maze as the brain may use this mechanism to regulate the activity of GABA which was seen in neuropathic animals reported by Saffapour *et al.* [33].

Increased expression due to agonist treatment can be explained that diazepam treatment initiated the increase in production of the receptors for binding causing increased inhibition leading to memory impairment.

Conclusion

NP induced plastic changes in the expression of GABA_A receptors in the brain and the spinal cord. GABA_A receptor agonist treatment has no ameliorative effect in NP as GABA signaling may have shifted to excitatory in nature.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- 1 Basbaum AI. Distinct neurochemical features of acute and persistent pain. *Proc Natl Acad Sci USA* 1999; 96:7739–7743.
- 2 Treede RD, Kenshalo DR, Gracely RH, Jones AK. The cortical representation of pain. *Pain* 1999; 79:105–111.
- 3 Yan Y, Li C, Zhou L, Ao L, Fang W, Li Y. Research progress of mechanisms and drug therapy for neuropathic pain. *Lif Sci* 2017; 190:68–77.
- 4 Jones RC, Lawson E, Backonja M. Managing neuropathic pain. *Med Clin N Am* 2016; 100:151–167.

- 5 Polgár E, Hughes DI, Riddell JS, Maxwell DJ, Puskár Z, Todd AJ. Selective loss of spinal GABAergic or glycinergic neurons is not necessary for the development of thermal hyperalgesia in the chronic constriction injury model of neuropathic pain. *Pain* 2003; 104:229–239.
- 6 Polgár E, Gray S, Riddell JS, Todd AJ. Lack of evidence for significant neuronal loss in laminae I–III of the spinal dorsal horn of the rat in the chronic constriction injury model. *Pain* 2004; 11:144–150.
- 7 Bennett GJ, Xie YK. A peripheral mononeuropathy in rat that produces disorders of pain sensation like those seen in man. *Pain* 1988; 33:87–107.
- 8 Moore KA, Kohno T, Karchewski LA, Scholz J, Baba H, Woolf CJ. Partial peripheral nerve injury promotes a selective loss of GABAergic inhibition in the superficial dorsal horn of the spinal cord. *J Neurosci* 2002; 22:6724–6731.
- 9 Coull JA, Boudreau D, Bachand K, Prescott SA, Nault F, Sik AD, *et al.* The trans-synaptic shift in anion gradient in spinal lamina I neurons as a mechanism of neuropathic pain. *Nature* 2003; 424:938–942.
- 10 Coull JA, Beggs S, Boudreau D, Boivin D, Tsuda M, Inoue K, *et al.* BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain. *Nature* 2005; 438:1017–1021.
- 11 Polgár E, Hughes DI, Arham AZ, Todd AJ. Loss of neurons from laminae I–III of the spinal dorsal horn is not required for the development of tactile allodynia in the spared nerve injury model of neuropathic pain. *J Neurosci* 2005; 25:6658–6666.
- 12 Scholz J, Broom DC, Youn DH, Mills CD, Kohno T, Suter MR, *et al.* Blocking caspase activity prevents trans-synaptic neuronal apoptosis and the loss of inhibition in lamina II of the dorsal horn after peripheral nerve injury. *J Neurosci* 2005; 25:7317–7323.
- 13 Polgár E, Sardella TCP, Tiong SYX, Locke S, Watanabe M, Todd AJ. Functional differences between neurochemically defined populations of inhibitory interneurons in the rat spinal dorsal horn. *Pain* 2013; 154:2606–2615.
- 14 Blom SM, Pfister JP, Santello M, Senn W, Nevian T. Nerve injury-induced neuropathic pain causes disinhibition of the anterior cingulate cortex. *J Neurosci* 2014; 34:5754–5764.
- 15 Abdulmajeed WI, Ibrahim RB, Ishola AO, Balogun WG, Cobham AE, Amin A. Amitriptyline and phenytoin prevent memory deficit in sciatic nerve ligation model of neuropathic pain. *J Bas Clin Physiol Pharmacol* 2015; 27:101–108.
- 16 Castro-Lopes JM, Tavares I, Coimbra A. GABA decreases in the spinal cord dorsal horn after peripheral neurectomy. *Brain Res* 1993; 620:287–291.
- 17 Ibuki T, Hama AT, Wang XT, Pappas GD, Sagen J. Loss of GABA-immunoreactivity in the spinal dorsal horn of rats with peripheral nerve injury and promotion of recovery by adrenal medullary grafts. *Neurosci* 1997; 76:845–858.
- 18 Eaton MJ, Plunkett JA, Karmally S, Martinez MA, Montanez K. Changes in GAD- and GABA-immunoreactivity in the spinal dorsal horn after peripheral nerve injury and promotion of recovery by a lumbar transplant of immortalized serotonergic precursors. *J Chem Neuroanat* 1998; 16:57–72.
- 19 Yau JL, Noble J, Hibberd C. Chronic treatment with the antidepressant amitriptyline prevents impairments in water maze learning in aging rats. *J Neurosci* 2002; 22:1436–1442.
- 20 Woolf CJ, Salter MW. Neuronal plasticity: increasing the gain in pain. *Science* 2000; 288:1765–1769.
- 21 Hu Y, Yang J, Hu Y, Wang Y, Li W. Amitriptyline rather than lornoxicam ameliorates neuropathic pain-induced deficits in abilities of spatial learning and memory. *Eur J Anaes* 2010; 27: 1628–1635.
- 22 Mutso AA, Radzicki D, Baliki MN, Huang L, Banisadr G, Centeno MV, *et al.* Abnormalities in hippocampal functioning with persistent pain. *J Neurosci* 2012; 32:5747–5756.
- 23 Kupers RC, Nyten DD, Castro-Costa M, Gytrels JM. A time course analysis of the changes in spontaneous and evoked behavior in a rat model of neuropathic pain. *Pain* 1992; 50:101–111.
- 24 Allen JW, Yaksh TL. Assessment of acute thermal nociception in laboratory animals. *Methods Mol Med* 2000; 99:11–23.
- 25 Honda K, Takano Y. Experimental methods in pain using neuropathic and inflammatory animal models. *Nipp Yakuri Zass* 2007; 130:39–44.
- 26 Ishola AO, Laoye BJ, Oyeleke DE, Bankole OO, Sirjao MU, Cobham AE, *et al.* Vitamin D3 receptor activation rescued corticostriatal neural activity and improved motor-cognitive function in D2R parkinsonian mice model. *JBSE* 2015; 8:601–615.
- 27 Chaplan SR, Guo HQ, Lee DH, Luo L, Liu C, Kuei C, *et al.* Neuronal hyperpolarization-activated pacemaker channels drive neuropathic pain. *J Neurosci* 2003; 23:1169–1178.
- 28 Ran R, Gu J, Fu J, Zhong H, Zhao Y, Gu YC, *et al.* The role of the GABA-A receptor of the adjacent intact dorsal root ganglion neurons in rats with neuropathic pain. *Acta Neurobiol Exp* 2014; 74:405–414.
- 29 Imlach WL, Bhola RF, Mohammadi SA, Christie MJ. Glycinergic dysfunction in a subpopulation of dorsal horn interneurons in a rat model of neuropathic pain. *Sci Rep* 2016; 6:37104.
- 30 Chen SL, Zang Y, Zheng WH, Wei XH, Liu XG. Inhibition of neuropathic pain by a single intraperitoneal injection of diazepam in the rat: Possible role of neurosteroids. *Chin J Physiol* 2016; 59:9–20.
- 31 Ford A, Castonguay A, Cottet M, Little JW, Chen Z, Symons-Liguori AM, *et al.* Engagement of the GABA to KCC2 signaling pathway contributes to the analgesic effects of A3AR agonists in neuropathic pain. *J Neurosci* 2015; 35:6057–6067.
- 32 Dzhalal VI, Talos DM, Sdrulla DA, Brumback AC, Mathews GC, Benke TA, *et al.* NKCC1 transporter facilitates seizures in the developing brain. *Nat Med* 2005; 11:1205–1213.
- 33 Saffapour S, Shaabani M, Naghdi N, Farahmandfar M, Janzad A, Nasirinezhad F. In vivo evaluation of the hippocampal glutamate and GABA and BDNF level associated with spatial memory performance in a rodent model of neuropathic pain. *Physiol Behav* 2017; 175: 97–103.
- 34 Cao XY, Xu H, Wu LJ, Li XY, Chen T, Zhuo M. Characterization of intrinsic properties of cingulate pyramidal neurons in adult mice after nerve injury. *Mol Pain* 2009; 5:73.
- 35 Khakpai F, Nasehi M, Haeri-Rohani A, Eidi A, Zarrindast MR. Septo-Hippocampo-Sepal loop and memory formation. *Bas Clin Neurosci* 2013; 4:5–23.
- 36 Palazzo E, Romano R, Luongo L, Bocella S, De Georger Gordano ME. MMP-13, an mGluR7-selective negative allosteric modulator, alleviates pain and normalizes affective cognitive behavior in neuropathic pain mice. *Pain* 2015; 156:1060–1073.
- 37 Siegel A, Sapru HN. *Essential neuroscience*; Chapter nine. 2nd ed. Baltimore, MD, USA: Wolters Kluwer Lippincott Williams and Wilkins; 2011.
- 38 Polgár E, Todd AJ. Tactile allodynia can occur in the spared nerve injury model in the rat without selective loss of GABA or GABA_A receptors from synapses in Laminae I–II of the ipsilateral spinal dorsal horn. *Neurosci* 2008; 156:193–202.
- 39 Qu C, King T, Okun A, Lai J, Fields HL, Porreca FX. Lesion of the rostral anterior cingulate cortex eliminates the aversiveness of spontaneous neuropathic pain following partial or complete axotomy. *Pain* 2008; 152:1641–1648.
- 40 Neugebauer V, Li W, Bird GC, Han JS. The amygdala and persistent pain. *Neuroscience* 2004; 10:221–234.
- 41 Saab CY. Pain-related changes in the brain: diagnostic and therapeutic potentials. *Trends Neurosci* 2012; 35:629–637.
- 42 Lefaucheur JP, Drouot X, Ménard-Lefaucheur I, Keravel Y, Nguyen JP. Motor cortex RTMS restores defective intracortical inhibition in chronic neuropathic pain. *Neurol* 2006; 67:1568–1574.