

**Original Article:****Awareness and Utilization of Cervical Cancer Screening among Women in an Urban Area in Southwestern Nigeria****Authors:**

**Deborah Tolulope Esan**, Senior Lecturer, Department of Nursing, College of Medicine and Health Sciences, Afe Babalola University Ado-Ekiti, Nigeria,

**Ayodeji Akinwande Fasoro**, Lecturer, Department of Public Health, College of Medicine and Health Sciences, Afe Babalola University Ado-Ekiti, Nigeria,

**Fayokemi Mary Olaiya**, Undergraduate student, Department of Nursing, College of Medicine and Health Sciences, Afe Babalola University Ado-Ekiti, Nigeria,

**Cecilia Bukola Bello**, Lecturer, Department of Nursing, College of Medicine and Health Sciences, Afe Babalola University Ado-Ekiti, Nigeria.

**Address for Correspondence**

**Ayodeji Akinwande Fasoro**,

Department of Public Health,

College of Medicine and Health Sciences,

Afe Babalola University,

P.M.B. 5464, Ado-Ekiti, Nigeria.

**E-mail:** akinfasoro@abuad.edu.ng.

**Citation**

Esan DT, Fasoro AA, Olaiya FM, Bello CB. Awareness and Utilization of Cervical Cancer Screening among Women in an Urban Area in Southwestern Nigeria. *Online J Health Allied Scs.* 2019;18(3):4. Available at URL: <https://www.ojhas.org/issue71/2019-3-4.html>

Submitted: Aug 8, 2019; Accepted: Oct 13, 2019; Published: Oct 30, 2019

**Abstract:** Introduction: Cancer of the cervix is the commonest cause of cancer death in women worldwide. This study assessed the awareness level and pattern of utilization of cervical cancer screening test among women of different professions in Ado-Ekiti, Nigeria. Methods: A descriptive cross-sectional design was used and respondents were selected from four different professions. Results: About 40% of respondents had heard of cervical cancer prior to the survey and only 40.5% were aware of cervical screening tests. Among those who were aware of the screening tests, 16% had ever utilized cervical cancer screening tests. Conclusion: This study established low cervical cancer awareness level and low utilization rate of cervical cancer screening among respondents. Therefore, there is need for sustained and concerted community health awareness most importantly at the grassroots in order to increase awareness and utilization rate of cervical cancer screening among the sampled populace.

**Key Words:** Cervical cancer, Awareness, Utilization, Screening

**Introduction:**

Cancer of cervix is the second most common cancer affecting women globally and also the most common cancer in developing countries (1). These cancerous lesions begin in the epithelia lining of the cervix. Most of the time, these cells do not transform suddenly into cancer cell but first of all appear as precancerous lesions which are detectable by screening. These lesions are detectable by Pap smear and sometimes by visual acetic staining (2).

Cervical cancer incidence has remained steady in developing countries with more and more cases seen as years rolled by, whereas in developed countries, cancer of cervix has declined significantly over the last two decades. A review showed that

an average of 19% of women in developing countries have been screened for cervical cancer as against 63% of women in the developed countries (3). The question is why are reproductive age women not going for screening in developing countries? Could this be attributed to misconceptions, misused idea and negative beliefs about cervical cancer or rather as a result of ignorance about existing/available screening services? Reports from previous study have revealed that women who are not screened are at a higher likelihood of having cervical cancer (4). This has made it paramount to continue to investigate reasons for low utilization rate of cervical cancer screening in Nigeria. Up-till date cervical screening face serious challenge in the developing countries, as lack of resources and lack of awareness limits its coverage (5). Women's knowledge is also implicated in screening uptake and this is necessary in order to facilitate uptake. Studies have revealed that low awareness level of cancer of cervix and its prevention is likely to limit uptake of screening (6,7,8). In a study conducted by among health professionals in Nigeria; the study revealed low uptake of cervical cancer screening. It is noteworthy that, in spite of adequate knowledge of this group of professionals on the importance of cervical cancer screening test, uptake was still low (9).

The problem posed by cervical cancer in Nigeria cannot be ignored, as any of the affected women report to hospital late when cancer prognosis is poor as a result of the cancer cells leaving their primary tumor site to infiltrate surrounding tissues and also distant organs in the body. This has reduced the survival rate of cancer related deaths in Nigeria. Moreover, there is high risk of developing cancer of the cervix among women of reproductive age and among sexually active teenagers, yet awareness of the disease is low. Uptake of

screening services for cervical cancer is done by only few who know about cervical cancer screening and its importance (10). It has been reported that women in Nigeria are not adequately informed about the disease, the predisposing factors, and are uncertain about the importance of the screening tests. Women also hold negative belief and perception about screening tests (11). This present work therefore assessed the awareness level and utilization pattern of cervical cancer screening among women in Ado-Ekiti, Ekiti State, Nigeria.

### Materials and Methods

The study employed a descriptive cross-sectional design. The study was conducted among women belonging to three different professions in the city of Ado-Ekiti. The participants included in this study were teachers, bankers and artisans. Teachers were recruited from Afe Babalola International Schools Ado-Ekiti and the Federal Polytechnic Staff School Ado-Ekiti. Afe Babalola International Schools comprise the primary and secondary schools. The Federal Polytechnic Staff School Ado-Ekiti has only a primary school. Female bankers from selected banks in Ado-Ekiti were recruited for the study. Female artisans (traders, tailors and hairdressers) were recruited for the study within Afe Babalola University community. The sampling technique used was purposive sampling technique. Sample size was computed using the Cochran equation in Fischer's formula,

$$n = \frac{Z^2 P(1-P)}{d^2}$$

Where, n is the desired sample size required, Z is the standard normal deviate corresponding to 95% confidence interval, P is the proportion of attribute (16%) obtained from a study conducted in South-Eastern part of Nigeria (12) and d is the margin of error of 5%. The estimated sample size required (n) was 206 participants.

The research instrument was an adapted semi-structured questionnaire. The questionnaire was self-administered to the participants to assess their awareness level and utilization of screening tests of cervical cancer. Relevant questions about the study were constructed in such a way that would provide relevant information about the study. The questionnaire was self-administered to 206 participants. The questionnaire contained three sections: section A contained the demographic profile of participants and it includes age, educational level, marital status, occupation, religion and ethnicity; section B contained questions on awareness level of respondents on cancer of cervix and its screening tests. Items included in this section assessed the respondents' awareness of the predisposing factors of cancer of the cervix, probable cause of cervical cancer, signs and symptoms of the disease, prevention and screening tests; and section C contained questions on utilization level/uptake of screening tests by respondents. After obtaining both written and verbal consent from the participants, the self-administered questionnaires were distributed to the participants at Afe Babalola University, Afe Babalola International Schools, Federal Polytechnic Staff School, and selected banks in Ado-Ekiti. The period of data collection lasted for three months (December 2017 – February 2018). Data were analyzed with SPSS version 20 using descriptive statistics.

### Results

Socio-demographic demographic profile of respondents is presented in Table 1. The mean age of the respondents was 29.7±5.8 years. Majority were between ages 21 – 30 years (54.9%), had tertiary education (71.8%), married (58.7%), Christians (84.0%), and belong to the Yoruba ethnic group (79.6%).

Table 1: Demographic Profile of respondents		
Item	Frequency (n=206)	Percentage (%)
<b>Age</b>		
Less than 21	8	3.9
21-30	113	54.9
31-40	77	37.3
41 and above	6	2.9
Missing data	2	1.0
Mean age ± Standard deviation	29.7±5.8	
<b>Level of education</b>		
Primary	4	1.9
Secondary	51	24.8
Tertiary	148	71.8
Missing data	3	1.5
<b>Marital status</b>		
Single	58	28.2
Married	121	58.7
Divorced	5	2.4
Separated	12	5.8
Missing data	10	4.9
<b>Religion</b>		
Christianity	173	84.0
Islam	30	14.5
Others	1	0.5
Missing Data	2	1.0
<b>Ethnicity</b>		
Yoruba	164	79.6
Hausa	14	6.8
Igbo	22	10.7
Others	6	2.9
<b>Occupation</b>		
Teaching	60	29.1
Banking	32	15.5
Hairdressing	40	19.4
Tailoring	50	24.3
Business/Trading	24	11.7

Approximately 40% of the women have heard of cancer of the cervix as shown in Table 2. Human papilloma virus (HPV) and having multiple sexual partners were the two most important risk factors identified by those who have ever heard of cervical cancer. Only 30.5% knew that there exists a vaccine which can help prevent cervical cancer.

Table 2: Awareness of Respondents on Cancer of cervix and cervical cancer screening test						
Item	Yes		No		Don't Know	
	N	%	n	%	n	%
Heard of cancer of the cervix before? (n=206)	82	39.8	124	60.2	0	0.0
Cancer of the cervix is a common cancer (n=82)	31	37.8	22	26.8	29	35.4
<b>Risk factors for cervical cancer include (n=82)</b>						
Having a relative that had the disease	31	37.8	24	29.3	27	32.9
Having HIV/AIDS	13	15.9	45	54.9	24	29.2
Having multiple sexual partner	41	50.0	18	22.0	23	28.0
Early marriage	22	26.8	29	35.4	31	37.8
Early initiation of sex	30	36.6	20	24.4	32	39.0
Human papilloma virus	48	58.5	11	13.4	23	28.1
Poverty	11	13.4	35	42.7	36	43.9
<b>The cause of cervical cancer (n=82)</b>						
Infection with HIV/AIDS	10	12.2	40	48.8	32	39.0
Infection with Human papilloma virus	53	64.6	7	8.6	22	26.8
Lack of personal hygiene	34	41.4	18	22.0	30	36.6
<b>Clinical manifestation of cervical cancer (n=82)</b>						
Abnormal vaginal bleeding	62	75.6	5	6.1	15	18.3
Bleeding after intercourse	30	36.6	24	29.3	28	34.1
Smelling vaginal discharge	45	54.9	14	17.1	23	28.0
Pain while urinating	28	34.2	26	31.7	28	34.1
Abdominal pain	27	32.9	23	28.1	32	39.0
<b>Prevention of cervical cancer (n=82)</b>						
Antibiotic use	19	23.2	28	34.1	35	42.7
Cervical screening	59	72.0	3	3.7	20	24.3
Immunization	25	30.5	18	22.0	39	47.5
Heard of cervical cancer screening before (n=206)	83	40.5	123	59.5	0	0.0
<b>Procedures done in cervical cancer screening (n=83)</b>						
Blood test	42	50.6	21	25.3	20	24.1
Urine test	20	24.1	36	43.4	27	32.5
X-ray	16	19.3	38	45.8	29	34.9
Ultrasound	23	27.7	30	36.2	30	36.1
Operation	16	19.3	38	45.8	29	34.9
Taking a sample from the cervix for examination/test	52	62.7	4	4.8	27	32.5
Visualizing after staining the cervix with acetic acid	40	48.2	4	4.8	39	47.0

Table 3 shows that only 16% have ever had cervical cancer screening and 54.6% of them voluntarily went for the screening. Most of them (54.5%) had the screening done in a government hospital.

Table 3: Utilization of cervical cancer screening test		
Item	n	%
<b>Ever gone for cervical cancer screening test</b>		
Yes	33	16.0
No	170	82.5
Missing data	3	1.5
<b>Reason(s) for cervical cancer screening</b>		
It was prescribed	8	24.2
It was voluntary	18	54.6
Suggested to me by my friend	7	21.2
<b>Where cervical cancer screening was done</b>		
Government hospital	18	54.5
Private hospital	15	45.5
<b>How many times cervical cancer screening had been done before</b>		
Once	16	7.8
Twice	11	5.3
Thrice	4	1.9
Four times	1	0.5
More than four times	1	0.5
Never	142	68.9
Missing data	31	15.1
<b>Previous date of cervical cancer screening test</b>		
This year	4	1.9
Last year	20	9.7
2 years ago	3	1.5
3-4 years ago	3	1.5
Never	143	69.4
Missing data	33	16.0
<b>Would you like to have regular screening for cervical cancer</b>		
Yes	49	23.8
No	134	65.0
Missing data	23	11.2
<b>Reason for not going for cervical cancer screening (n=104)</b>		
I may not keep to appointments	58	55.8
I don't like it, the procedure is discomforting	29	27.9
I don't like male staff attending to me	12	11.5
Others	5	4.8
<b>Ever advised someone to go for cervical cancer screening</b>		
Yes	65	31.6
No	138	67.0
Missing data	3	1.4
<b>Will you encourage your friends and relatives to go for screening</b>		
Yes	89	43.2
No	110	53.4
Missing data	7	3.4

### Discussion

Respondents mean age was 29.7±5.8 years and the ages ranged from 19 – 48years. This is contrary to the results obtained in a study conducted by Arulogun and Maxwell (13) in Ibadan, Oyo State Nigeria, where the mean age was 33.9 years. The tribal distribution showed that 79.6%, 10.7% and 6.8% were Yoruba, Hausa and Igbo respectively. This is in contrast with a study in Sokoto, Nigeria which reported the tribal distribution of 35.9% Hausa, 21.8% Yoruba and 21.8% Igbo

(14). The reason for this disparity is because their research was conducted in the north western part of Nigeria where majority of the population are Hausas, and this present research was conducted in Ekiti State, Southwest, Nigeria where majority are Yorubas. The age distribution, marital status, and educational level of women in this study is similar to the reports of other studies carried out in Nigeria (15-17).

#### **Cervical cancer screening awareness among women of different profession**

About 40% of respondents had heard of cervical cancer prior to the survey. This is contrary to the result reported by Oche and colleagues (14) where almost all (98.6%) the respondents had heard of cancer of the cervix. The reason for this contrast is because; their study was conducted among female health workers whereas this present research was conducted among women of different professions, which included artisans (tailors, hairdressers and business owners), teachers and bankers. About 58.5% identified HPV and 50% multiple sex partners as predisposing factors for cancer of the cervix. Higher percentage was obtained from a research conducted by Oche and colleagues (14) where 85.5% and 87.7% were aware that cancer of the cervix has a link with HPV and multiple sexual partners respectively.

More than half (59.5%) of the respondents were not aware of cervical cancer screening test, while a lesser percentage (40.5%) who have heard of the screening. These findings are a bit different from what was obtained in a study conducted by Ogunbode and Ayinde (15) where only 19.7% of market women in Ibadan knew about screening test for cervical cancer. The difference in awareness level may be due to the fact that their study was conducted amidst market women whereas this present study was conducted among women of different professions. However, the findings of our study are similar to what was obtained in a study conducted among rural women in Osun state where awareness level was 39.2% (18). This study finding was also in contrast to the findings by Ahmed and colleagues (17) among market women in Zaria where 68.9% had knowledge of the screening services. Aboyeji and colleagues (19) conducted a study in Ilorin, Nigeria where awareness about cervical cancer screening was 69.8% and 87% in another study among female health professionals (14). These findings however cannot be compared because the studies were conducted among different study populations with different academic backgrounds and qualifications.

#### **Utilization of screening test of cervical cancer among respondents**

The findings from this study showed that only few (16%) of the respondents' ever utilized cancer of the cervix screening tests. This finding is however lower to what was obtained in another study (17), where about one third (32.7%) of the women utilized cervical cancer screening services. The report obtained in this study is however higher than what was obtained in a study conducted in Enugu, South-Eastern, Nigeria (20) where only about 2.8% of the women had ever utilized cervical cancer screening services. Furthermore, a study conducted in Owerri also reported lower usage (7.1%) of the screening (21). Our study reveals that 7.8% of the respondents had gone for cervical screening once and only 0.5% had gone more than four times. Similar to this, as noticed by previous works, utilization of cervical cancer screening was similarly poor.

This study has a few limitations worth mentioning. The participants recruited for this study were selected through a non-random sampling procedure. Hence, there could have been selection bias and the result of this study cannot be generalized to the entire women population in the town, state or country. The cross-sectional study design employed cannot establish a cause-effect relationship. Also, it is a

questionnaire-based study; so, reporting bias cannot be ignored.

#### **Conclusion**

There was a low awareness level of cervical cancer screening test among respondents in the study population and utilization pattern of these services was also poor. Only a few of the women are knowledgeable about cervical cancer with only a few also aware of the screening services. A lot still has to be done in the area of awareness creation and initiating cultural, acceptable, feasible and cost effective screening programs. We suggest that these screening services should be made available free of charge at all primary health care centers in the State, if the fight against the disease is to be won.

**Conflict of Interest:** All authors declare no conflict of interest.

**Funding:** No funding was received for the conduct of this research

#### **References**

1. Urasa M, Darj E. Knowledge of cervical cancer and screening practices of nurses at a regional hospital in Tanzania. *Afr Health Sci*. 2011; 11(1):48-57.
2. Modibbo FI, Dareng E, Bamisaye P, Jedy-Agba E, Adewole A, Oyeneyin L, Olaniyan O, Adebamowo C. Qualitative study of barriers to cervical cancer screening among Nigerian women. *BMJ Open*. 2016; 6(1):e008533.
3. Ncube B, Bey A, Knight J, Bessler P, Jolly PE. Factors associated with the uptake of cervical cancer screening among women in Portland, Jamaica. *North American Journal of Medical Sciences*. 2015; 7(3):104.
4. Oscarsson MG, Benzein EG, Wijma BE. Reasons for non-attendance at cervical screening as reported by non-attendees in Sweden. *Journal of Psychosomatic Obstetrics & Gynecology*. 2008; 29(1):23-31.
5. Bansil P, Wittet S, Lim JL, Winkler JL, Paul P, Jeronimo J. Acceptability of self-collection sampling for HPV-DNA testing in low-resource settings: a mixed methods approach. *BMC Public Health*. 2014; 14(1):596.
6. Abotchie PN, Shokar NK. Cervical cancer screening among college students in Ghana: knowledge and health beliefs. *International Journal of Gynecological Cancer*. 2009; 19(3):412.
7. Liao CC, Wang HY, Lin RS, Hsieh CY, Sung FC. Addressing Taiwan's high incidence of cervical cancer: factors associated with the Nation's low compliance with Papanicolaou screening in Taiwan. *Public Health*. 2006; 120(12):1170-6.
8. Hummeida M, Elrasheed T, Burhan A. Cervical cancer prevention in Sudan. Barriers and missed opportunities. *International Journal of Gynecology & Obstetrics*. 2009; 107(S2).
9. Chukwuali LI, Onuigbo WI, Mgbor NC. Cervical Cancer Screening in Enugu, Nigeria. *Tropical Journal of Obstetrics and Gynaecology*. 2003; 20(2):109-12.
10. WHO. WHO/ICO Information Centre on HPV and Cervical Cancer (HPV Information Centre). Human Papillomavirus and Related Cancers in South Africa. Summary Report 2010. Available at: [www.who.int/hpvcentre](http://www.who.int/hpvcentre)
11. Demirtas B, Acikgoz I. Promoting Attendance at Cervical Cancer Screening: Understanding the Relationship with Turkish Women's Health Beliefs. *Asian Pacific Journal of Cancer Prevention*. 2013; 14(1):333-40.
12. Feyi-Waboso PA, Kamanu C, Aluka C. Awareness and risk factors for cervical cancer among women in Aba,

- south-eastern Nigeria. *Tropical Journal of Obstetrics and Gynaecology*. 2005; 22:25–6.
13. Arulogun OS, Maxwell OO. Perception and utilization of cervical cancer screening services among female nurses in University College Hospital, Ibadan, Nigeria. *Pan African Medical Journal*. 2012; 11(1).
  14. Oche MO, Kaoje AU, Gana G, Ango JT. Cancer of the cervix and cervical screening: Current knowledge, attitude and practices of female health workers in Sokoto, Nigeria. *International Journal*. 2013; 5(3):106-9.
  15. Ogunbode OO, Ayinde OA. Awareness of cervical cancer and screening in a Nigerian female market population. *Annals of African Medicine*. 2005;4:160–3
  16. Nwozor CM, Oragudosi AL. Awareness and uptake of cervical cancer screening among women in Onitsha, South-East, Nigeria. *Greener Journal of Medical Sciences*. 2013; 3(8):283-8.
  17. Ahmed SA, Sabitu K, Idris SH, Ahmed R. Knowledge, attitude and practice of cervical cancer screening among market women in Zaria, Nigeria. *Nigerian Medical Journal: Journal of the Nigeria Medical Association*. 2013; 54(5):316.
  18. Ojo MI, Olowokere AE. Cervical cancer screening knowledge and utilisation among rural women in Ife East local government area of Osun State in Nigeria. *Africa Journal of Nursing and Midwifery*. 2014; 16(1):145-57.
  19. Aboyeji PA, Ijaiya MD, Jimoh AG. Knowledge, attitude and practice of cervical smear as a screening procedure for cervical cancer in Ilorin, Nigeria. *Tropical Journal of Obstetrics and Gynaecology*. 2004; 21(2):114-7.
  20. Ingwu JA. Knowledge and Screening Practices of Cervical Cancer among Pregnant Women Attending Antenatal Clinic in Tertiary Hospitals in Enugu, South-Eastern Nigeria. *Journal of Cancer and Tumor International*. 2016; 4(2):1-9.
  21. Ezem BU. Awareness and uptake of cervical cancer screening in Owerri, South-Eastern Nigeria. *Annals of African Medicine*. 2007; 6(3):94.