

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/328687462>

Evaluation of Goitrogenic Content of Common Vegetables in South West Nigeria

Article in *Journal of Food Science* · August 2018

DOI: 10.9734/AFSJ/2018/42750

CITATIONS

2

READS

463

5 authors, including:



Afees Adebayo Oladejo

Nnamdi Azikiwe University, Awka


17 PUBLICATIONS 9 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Evaluation of Goitrogenic Content of Common Vegetables in South West Nigeria [View project](#)



Evaluation of Goitrogenic Content of Common Vegetables in South West Nigeria

**Oladejo, Afees Adebayo^{1*}, Okesola, Mary Abiola²,
Oyerinde, Adebawale Samuel³, Jaiyesimi, Kikelomo Folake⁴
and Kolawole, Jesumiluyi Augustine⁵**

¹*Department of Chemical Sciences and Biochemistry, Afe Babalola University, Ado-Ekiti, Ekiti State, Nigeria.*

²*Department of Chemical Sciences, Covenant University, Ogun State, Nigeria.*

³*Department of Biochemistry, Ekiti State University, Ado Ekiti, Nigeria.*

⁴*Biochemistry Unit, Department of Chemical Sciences, Afe Babalola University, Ado-Ekiti, Ekiti State, Nigeria.*

⁵*Department of Chemical Science, Federal Polytechnic Bida, Niger State, Nigeria.*

Authors' contributions

"This work was carried out in collaboration between all authors. Author OAA designed the study, performed the statistical analysis and wrote the protocol. Author OMA wrote the first draft of the manuscript. Authors OAS and JKF managed the analyses of the study. Author KJA managed the literature searches. All authors read and approved the final manuscript."

Article Information

DOI: 10.9734/AFSJ/2018/42750

Editor(s):

(1) Uttara Singh, Assistant Professor, Department of Foods and Nutrition, Govt. Home Science College, Panjab University, India.

Reviewers:

(1) J. U. Nwamarah, University of Nigeria, Nsukka, Nigeria.

(2) Dodo Juliet Dingtsen, University of Jos, Nigeria.

(3) Ndomou Mathieu, University of Douala, Cameroon.

Complete Peer review History: <http://www.sciencedomain.org/review-history/25941>

Original Research Article

Received 7th May 2018
Accepted 11th July 2018
Published 18th August 2018

ABSTRACT

Endemic goiter and associated iodine deficiency disorders (IDD) are prevalent in south west Nigeria. The present study was undertaken to identify the role of dietary goitrogen in the etiology of endemic goiter. Perchlorate, fluoride, nitrate, bromide, chloride, phosphate and cyanide content of common vegetables viz., cabbage, African Eggplant, Giant pigweed, Scent leaf, Amaranth, Tree Spinach, Black nightshade, consumed by the population of the region were measured. All the dietary goitrogen content in the investigated vegetables were found to be within the Acceptable

*Corresponding author: E-mail: oludeledamilola@gmail.com, adebayoo997@gmail.com;

Daily Intake and this observation suggests that in addition to iodine deficiency dietary intake of a cyanogenic plant, the combination of chemicals may play some role for the persistence of endemic goiter in Southwest Nigeria.

Keywords: Goiter; vegetables; goitrogens.

1. INTRODUCTION

Vegetables often mean the part of a plant that is edible other than a sweet fruit or seed. It thus typically implies the leaf, stem or root of a plant [1]. Vegetables play a vital role in human nutrition; they serve as the most rapid and lowest cost source of fibers, minerals and vitamins to the majority of people in developing countries, where they are often eaten in relatively small amounts as a side dish or relish with the staple foods [2]. The broad variation in taste, texture and colors of diverse vegetables is an exciting additional touch to the meals [3]. The use of many plants for food is often limited by the composition of goitrogenic substances in them as they remain hazardous to both man and animals [4]. Hence, vegetables are widely cultivated and consumed for their flavors and nutritional benefits [5]. Vegetables may also contain goitrogenic substances, which are detrimental to nutrition and health [6].

Goitrogens are substances which cause an enlargement of the thyroid gland, a condition known as goiter. They may act directly or indirectly on the gland by altering the regulatory mechanisms of the gland and peripheral metabolism and excretion of triiodothyronine (T3)/ thyroxine (T4) [7]. The thyroid gland is one of the sensitive endocrine hormone producing glands that enhance protein synthesis and oxygen utilization, which in turn, influence the basal metabolic rate (BMR). The thyroid accomplishes the task of metabolic regulation through the secretion of three important hormones, T4 (thyroxine) and T3 (triiodothyronine) [8,9], as well as calcitonin [10], which is important for calcium metabolism in the blood plasma. When there is difficulty in making thyroid hormone, the thyroid gland enlarges as a way of trying to compensate for this inadequate production of the hormone. The level of thyroid hormone production is determined by the level of thyroid stimulating hormone (TSH) released from the pituitary gland; and by the availability of iodine and tyrosine. A number of compounds, however, have the ability to prevent the synthesis of these thyroid hormones. These compounds include perchlorate, fluoride, nitrate, bromide,

chloride, phosphate and cyanide. These compounds are generally gotten from food and water [7]. Association between goitrogens and goiter has been studied in detail [11]. Cassava (*Manihot utilissima*) [12,13], Cabbage (*Brassica oleracea* L.) [14,15] and Pearl Millet (*Pennisetum americanum*) [16,17,18] are few common examples of foods rich in goitrogens, which have been reviewed extensively. In spite of numerous reports on goitrogen contents in food, there is a dearth of literature report on the goitrogen levels in leafy vegetables. Hence, Investigation of the levels of goitrogens in vegetables is thus needed to discern the contribution of the vegetable to nutrition and health. This study thus evaluates the composition of goitrogens in some Nigerian leafy vegetables.

2. MATERIALS AND METHODS

2.1 Plant Materials

Samples of fresh vegetable leaves were collected from a farmland in Ado Ekiti, Ekiti State, Nigeria. The geographical coordinate of the farmland is Latitude: 7°37.3974' N; Longitude: 5°13.2522' E. Authentication of the plants was carried out at the Department of Plant Science and Forestry, Ekiti State University, Nigeria. The leaves were separated from the stalk, rinsed with distilled water and were oven-dried at 40°C. The dried samples were grounded into fine powder and sieved through 2.0 mm mesh prior to analysis.

2.2 Goitrogen Analysis

The Goitrogenic component of the samples was determined using the method as described by Tel and Rao [19]. Briefly, 20 g of each blended sample was mixed with 20 ml of distilled water and filtered through Whatman no. 2 filter paper, and the filtrate was passed through a glass column fitted with a tape and filled with activated alumina, in order to separate the green colour (chlorophyll) and get a transparent solution. The transparent solution, free from turbidity was used for the determination of the goitrogenic parameters.

Table 1. Selected leafy vegetables identified in Ekiti State

Scientific name	Common name	Vernacular name
<i>Brassica oleracea</i>	Cabbage	'Araasa'
<i>Solanum macrocarpon</i>	African Eggplant	'Igbagba'
<i>Trianthema portulacastrum</i>	Giant pigweed	'Esisan'
<i>Ocimum gratissimum</i>	Scent leaf	'Efinrin'
<i>Amaranthus cruentus</i>	Amaranth	'Arowojeja'
<i>Amaranthus viridis</i>	Amaranth	'Tete Adayeba'
<i>Cnidoscolus aconitifolius</i>	Tree Spinach	'Iyana-ipaja'
<i>Solanum nigrum</i>	Black nightshade	'Odu'

Table 2. Goitrogen composition of Nigerian leafy vegetables

Leafy vegetable	Nitrate (ppm)	Perchlorate (ppm)	Fluoride (ppm)	Bromide (ppm)	Chloride (ppm)	Phosphate (ppm)	Cyanide (ppm)
<i>Brassica oleracea</i>	0.15	0.08	2.15	0.08	0.54	0.04	0.45
<i>Solanum macrocarpon</i>	0.54	0.11	3.21	0.12	0.80	0.06	0.60
<i>Ocimum gratissimum</i>	0.20	0.05	1.48	0.04	1.17	0.12	0.32
<i>Amaranthus cruentus</i>	0.48	0.08	2.14	0.08	0.54	0.04	0.21
<i>Amaranthus viridis</i>	0.18	0.04	1.22	0.04	0.30	0.02	0.37
<i>Solanum nigrum</i>	0.27	0.09	2.48	0.09	0.62	0.05	0.53
<i>Cnidoscolus aconitifolius</i>	0.28	0.06	1.81	0.07	0.45	0.04	0.44
<i>Trianthema portulacastrum</i>	0.28	0.13	3.70	0.14	0.92	0.07	0.71

3. RESULTS AND DISCUSSION

The results of the Goitrogen analysis of the leafy vegetables are represented in Table 2. Nitrate, a naturally occurring compound and a vital component of vegetables because of its potential to a mass is formed naturally in living and decomposing plants and animals, including humans [20,21,22,23]. Nitrate by itself is relatively nontoxic, but its metabolites, nitrite, nitric oxide and N-nitroso compounds, make nitrate of regulatory importance because of their potentially harmful health implications. Nitrate majorly gets into the human body exogenously from vegetables, water, and other foods, but is also formed to a minute extent endogenously [21,22]. Nitrate is present in the investigated vegetables at different concentrations. The toxicity of Nitrate is due to its conversion to nitrite and its ability to react to form N-nitroso compounds whose effects have been identified to range from methemoglobin formation, hyperplasia of the zona glomerulosa of the adrenal cortex and gastric neoplasia [24]. The

safety of this compound is therefore dependent on its availability in food. However, the levels of nitrate in all the investigated vegetables are within the limit of the Acceptable Daily Intake (ADI) of 0-4000ppm/ day [25].

Perchlorate is a contaminant present in the environment naturally and as a result of human activity [26]. Hence, the use of natural fertilizers and perchlorate contaminated irrigation water may lead to considerable concentrations in leafy vegetables. Perchlorate is often found as the anion component of salt usually associated with cations such as ammonium, sodium or potassium. Perchlorate acts as a goitrogenic substance because it competitively blocks iodide from entering the thyroid due to its similarity in ionic radius to iodide by an effect on the Na^+/I^- symporter thus preventing the further synthesis of thyroid hormone, thereby inhibiting growth and development [27]. Severe iodine deficiency as a result of insufficient iodine intake or sustained exposure to goitrogenic substances such as perchlorate at levels that induce depletion of the

thyroid hormone stores can result in hypothyroidism. However, a mild to moderate iodine deficiency can lead to the development of toxic multinodular goiter and can result in hyperthyroidism. Besides its potential to cause endocrine system and reproductive problems, perchlorate is considered to be carcinogenic to humans [28]. The widespread presence of perchlorate in vegetables and its toxicological properties has however made perchlorate an emerging chemical of concern. From this study, the levels of perchlorate in the investigated vegetables are within the tolerable daily intake (TDI) of 0-0.3 ppm per day [29]. Meanwhile, previous research on perchlorate and possible thyroid-related health effects has paid slim emphasis on the other common environmental Sodium-iodide co-transporter inhibitors, thiocyanate and nitrate. The focus on perchlorate grow in part due to its relative potency as a Sodium-iodide co-transporter inhibitor is 10-200 times that of thiocyanate, and nitrate respectively on a molar basis [30,31]. However, based on average daily intake of perchlorate equivalents of nitrates and thiocyanates, a person's exposure to both thiocyanate and nitrate from food account for a larger proportion of iodine uptake inhibition than does perchlorate exposure [32]. Moreover, *in vitro* studies of NIS indicate that perchlorate, nitrate, and thiocyanate act synergistically to inhibit iodide uptake [31]. Therefore, it seems very vital to study the combination of these chemicals.

Bromide and chloride have been reported to have similar biological behavior on the thyroid gland. High bromide intake has been shown to decrease the amount of iodide accumulated in the thyroid [33], thus influencing their iodine metabolism by reducing the accumulation of iodine in the thyroid gland and skin as well as by a rise in iodide excretion by the kidneys, thereby exerting goitrogenic effects. Very high consumption of bromide has been linked to a short biological half-life of iodine in the thyroid of rats [33]. However, the concentration of these compounds in the investigated vegetables is far below that which can result in acute and chronic toxicity [34]. Phosphate is widely distributed in the body and is involved in cell signaling, energy metabolism, nucleic acid synthesis as well as maintenance of acid-base balance. Excess phosphate in the blood can lead to calcium deposits in the soft tissues of the body; a condition evident in the patient with reduced kidney function. The level of phosphate observed

in the investigated vegetables is however far from that which can be toxic to humans [28].

4. CONCLUSION

The goitrogenic potential of a plant or food depends upon the amount of active goitrogen present in it. The consumption of vegetables, though rich in nutrients could be limited and dangerous to health if their goitrogenic content is high beyond the acceptable limit. However, the results of this study showed that the goitrogenic substances in Nigerian vegetable are in safe amount. Hence, consumption of these vegetables is beneficial to health.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Omale J, Ugwu CE. Comparative studies on the protein and mineral composition of some selected Nigerian vegetables. *African Journal of Food Science*. 2011; 5(1):22-25.
2. Iyaka YA. Concentration of Cu and Zinc in some fruits and vegetables commonly available in North-Central zone of Nigeria. *Electronic Journal of Environmental, Agricultural and Food Chemistry*. 2007; 6(6):2150-2154.
3. Akindahunsi AA, Salawu SO. Phytochemical screening and nutrient-antinutrient composition of selected tropical green leafy vegetables. *African Journal of Biotechnology*. 2005;4(6):497-501.
4. Kubmarawa D, Andenyand IFH, Magomya AM. Amino acid of two non-conventional leafy vegetables: *Sesamum* and *Balanites aegyptiaca* L. *Africa Journal of Biotechnology*. 2008;7(19):3502-3504.
5. Yahaya AI, Suleiman I, Rahmatallah AA, Bello UB. Nutrient Content of Selected Edible Leafy Vegetables. *American Journal of Applied Chemistry*. 2014;2(3):42-45.
6. Yuwei L, Weihua X, Xiaoxiao J, Bo Z, Qian W, Yijian H. The impact of processing on phytic acid, *in vitro* soluble zinc and Phy/Zn molar ratio of faba bean (*Vicia faba* L.) *International Food Research Journal*. 2013;20(3):1285-1291.

7. Gaitan E. Goitrogens in food and water. *Annual Review of Nutrition*. 1990;10:21-39.
8. Joseph CND. Phytotherapeutic support of thyroid function. *Nutrition News*. 2007;8: 1-4.
9. Yilmaz S, Ozan S, Benzer F, Canatan H. Oxidative damage and antioxidant enzyme activities in experimental hypothyroidism. *Cell Biochemistry Function*. 2003;21:325-330.
10. Ane GM, Silvia SP, Francisco GG, Juan CG. The incidence and prevalence of thyroid dysfunction in Europe: A meta-analysis. *Archives of Internal Medicine*. 2013;160:526-34.
11. Chudasama RK, Verma PB, Mahajan RG. Iodine nutritional status and goiter prevalence in 6-12 years primary school children of Saurashtra region, India. *World Journal of Pediatrics*. 2010;6(3):233-237.
12. Akinpelu AO, Amangbo EF, Olojede AO, Oyekale AS. Health implications of cassava production and consumption. *Journal of Agriculture and Social Research (JASR)*. 2011;11(1):1-3.
13. Sajid L, Joachim M. Potential of cassava leaves in human nutrition: A review. *Trends in Food Science and Technology*. 2015;442(2).
14. Latham MC. A goitre survey in Ukinga, Tanzania. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 1965;59(3):342-348.
15. Fernando R, Pinto MDP, Pathmeswaran A. Goitrogenic Food and prevalence of Goitre in Sri Lanka. *Int. J. Internal Medicine*. 2012;1(3):17-20.
16. Eduardo G, Raymond HL, Robert DR, Sidney HI, Robert CC, Jim L, Edward FM, John H, Ken K. Antithyroid and Goitrogenic Effects of Millet: Role of C-Glycosyl flavones. *The Journal of Clinical Endocrinology and Metabolism*. 2009; 68(4):4-7.
17. Elnour A, Hambraeus L, Eltom M, Dramaix M, Bourdoux P. Endemic goiter with iodine sufficiency: A possible role for the consumption of pearl millet in the etiology of endemic goiter. *American Journal of Clinical Nutrition*. 2000;71(1):59-66.
18. Goldie1 JB, Atawodi SE, Isah HS, Berezi EP. Effect of Combined Millet and Cabbage Balanced Diet on Thyroid Function in Albino Rats. *Journal of Biology Agriculture and Healthcare*. 2014;4(20): 2224-3208.
19. Tel AD, Rao PV. Automated and Semi automated method of leafy vegetables analysis. Manual series. International Institute of Tropical Agriculture. Ibadan, Nigeria; 1982.
20. Mensinga TT, Speijers GJ, Meulenbelt J. Health implications of exposure to other environmental sodium/iodide symporter inhibitors: Potential thyroid-related health effects pollution in aquatic ecosystems: A global assessment. *Environment International*. 2003;32:831-849.
21. Lundberg JO, Weitzberg E, Cole JA, Benjamin N. Nitrate, bacteria and human health. *Nature Reviews Microbiology*. 2004;2:593-602.
22. Lundberg JO, Weitzberg E, Gladwin MT. The nitrate-nitrite-nitric oxide pathway in physiology and therapeutics. *Nature Reviews Drug Discovery*. 2008;7:156-167.
23. Camargo JA, Alonso A. Ecological and toxicological effects of inorganic nitrogen pollution in aquatic ecosystems: A global assessment. *Environmental International*. 2006;32:831-849.
24. Lundberg JO, Weitzberg E. NO generation from nitrite and its role in vascular control. *Arteriosclerosis, Thrombosis, and Vascular Biology Journal*. 2005;25:915-922.
25. EC (European Commission). Opinion on nitrate and nitrite. *Reports of the Scientific*. 1992;5-11.
26. Environmental Protection Agency (EPA). Federal Facilities Restoration and Reuse Office (FFRRO). Perchlorate treatment technology update- Federal Facilities Forum Issue Paper. 2005;542.
27. Wolff J. Perchlorate and the thyroid gland. *Pharmacological Reviews*. 1998;50(1):89-105.
28. Environmental Protection Agency (EPA). Air Quality Modeling Technical Support Document for the 2007 Fine Scale Modeling Platform. *Environmental nitrogenous compounds. Toxicological Reviews*. 2013;22:41-51.
29. European Food Safety Authority (EFSA). Scientific opinion on the risk to public health related to the presence of perchlorate in food, in particular fruits and vegetables. *European Food and Ankle Surgery (EFAS) Journal*. 2014;12(10): 3869.
30. Greer MA, Stott AK, Milne KA. Effects of thiocyanate, perchlorate and other anions on health. *Nature Reviews Microbiology*. 1966;2:593-602.

31. Tonacchera M, Pinchera A, Dimida A, Ferrarini E, Agretti P, Vitti P. Relative thyroidal iodine metabolism. *Endocrinology*. 2004;79:237-247.
32. De Groef B, Decallonne BR, Van der GS, Darras VM, Bouillon R. Perchlorate versus other environmental sodium/iodide symporter inhibitors: Potential thyroid-related health effects. *European Journal of Endocrinology*. 2006;155:17-25.
33. Stanislav P. Bromide interference with iodine metabolism. *Comprehensive Handbook of Iodine*. 2009;587-595.
34. Biller J. Interface of neurology and internal medicine (illustrated ed.). Lippincott Williams and Wilkins. 2007;939.

© 2018 Adebayo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/25941>